arXiv:1609.06535v1 [hep-th] 21 Sep 2016

Holographic phase transition probed by non-local observables

Xiao-Xiong Zeng[1,2*], Li-Fang Li[3†]

[1] School of Material Science and Engineering, Chongqing Jiaotong University, Chongqing 400074, China

[2]State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

[3] Center for Space Science and Applied Research, Chinese Academy of Sciences, Beijing 100190, China

September 22, 2016

Abstract

From the viewpoint of holography, the phase structure of a 5-dimensional Reissner-Nordström-AdS black hole is probed by the two point correlation function, Wilson loop, and entanglement entropy. As the case of thermal entropy, we find for all the probes, the black hole undergos a Hawking-Page phase transition, a first order phase transition and a second order phase transition successively before it reaches to a stable phase. In addition, for these probes, we find the equal area law for the first order phase transition is valid always and the critical exponent of the heat capacity for the second order phase transition coincides with that of the mean field theory regardless of the size of the boundary region.

*E-mail: xxzeng@itp.ac.cn

†E-mail: lilf@itp.ac.cn

Contents

1 Introduction **2**

2 Thermodynamic phase transition of the 5-dimensional Reissner-Nordström-AdS black hole **4**

3 Phase transition in the framework of holography **8**
3.1 Phase transition of two point correlation function 8
3.2 Phase transition of Wilson loop 12
3.3 Phase transition of entanglement entropy 15

4 Concluding remarks **19**

1 Introduction

Phase transition is a ubiquitous phenomenon for garden-variety thermodynamic systems. Due to the pioneering work by Hawking [1, 2], a black hole is also a thermodynamic system. Such a fact is further supported by AdS/CFT correspondence [3, 4, 5], where a black hole in the AdS bulk is dual to a thermal system without gravity. So one can naturally expect a black hole can also undertake some interesting phase transitions as the general thermodynamic system. Actually it has been shown that a charged AdS black hole undergos a Hawking-Page phase transition [6, 7], which is interpreted as the confinement/deconfinement phase transition in the dual gauge field theory [8], and a Van der Waals-like phase transition before it reaches the stable state [9]. The Hawking-Page phase transition implies that the thermal AdS is unstable and it will transit to the stable Schwarzschild AdS black hole lastly. The Van der Waals-like phase transition has been observed till now in many circumstantces. The first observation was contributed by [9] in the $T-S$ plane. Specifically speaking, in a fixed charge ensemble, for a black hole endowed with small charge, there is an unstable black hole interpolating between the stable small hole and stable large hole, and the small stable hole will undertake a first order phase transition to the large stable hole as the temperature of the black hole reaches a critical temperature. As the charge increases to the critical charge, the small hole and the large hole merge into one and squeeze out the unstable phase so that an inflection point emerges and the phase transition is second order. When the charge exceeds the critical charge, the black hole is always stable. Recently in the extended phase space, where the negative cosmological constant is treated as the pressure while its conjugate acts as the thermodynamical volume, the Van der Waals-like phase transition has also been observed in the $P-V$ plane [10, 11, 12, 13, 14, 15, 16]. In addition, it was shown in [17] that the Van der Waals-like phase transition also shows up in the $Q-\Phi$ plane. Particularly, in the Gauss-Bonnet gravity, it is found that the Gauss-Bonnet coupling parameter α also affects the phase structure of the space time, and in the $T-\alpha$ plane, a 5-dimensional neutral Gauss-Bonnet black hole also demonstrates the Van der Waals-like phase transition [18].

In this paper, we intend to probe the Hawking-Page phase transition and Van der Waals-like phase transition appeared in a 5-dimensional Reissner-Nordström-AdS black hole by the geodesic length, minimal area surface, and minimal surface area in the bulk, which are dual to the non-local observables on the boundary theory by holography, namely the two point correlation function, Wilson loop, and entanglement entropy, individually[1]. In fact, there have been some similar works to probe the phase structure by holographic entanglement entropy. In [26], the phase structure of entanglement entropy is studied in the $T-S$ plane for both a fixed charge ensemble and a fixed chemical potential ensemble, and it is found that the phase structure of entanglement entropy is similar to that of the thermal entropy. In particular, the entanglement entropy is found to demonstrate the same second order phase transition at the critical point as the thermal entropy. Soon after, it is found that the entanglement entropy can also probe the Van der Waals-like phase transition in the $P-V$ plane [27]. In [28], Nguyen has investigated exclusively the equal area law of holographic entanglement entropy and found that the equal area law holds regardless of the size of the entangling region. Very recently [29] investigated entanglement entropy for a quantum system with infinite volume, their result showed that the entanglement entropy also exhibits the same Van der Waals-like phase transition as the thermal entropy. They also checked the equal area law and obtained the critical exponent of the heat capacity near the critical point.

In this paper, we will further investigate whether one can probe the phase structure by two point correlation function and Wilson loop besides the entanglement entropy. We intend to explore whether they exhibit the similar Van der Waals-like phase transition as the entanglement entropy and thermal entropy. In addition, we also want to check whether these non-local observables can probe the Hawking-Page phase transition between the AdS black hole and thermal gas so that we can get a complete picture about the phase transition of the black holes in the framework of holography.

This paper is organized as follows. In the next section, we will discuss the thermal entropy phase transition of a 5-dimensional Reissner-Nordström-AdS black hole in the $T-S$ plane in a fixed charge ensemble. Then in Section 3, we will probe these phase transitions by geodesic length, Wilson loop, and holographic entanglement entropy individually. In each subsection, the equal area law is checked and the critical exponent of the heat capacity is obtained for different sizes of the boundary region. The last section is devoted to discussions and conclusions.

[1]Recently these non-local observables have been used to probe the non-equilibrium thermalization process, and it has been found that all of them have the same effect [19, 20, 21, 22, 23, 24, 25].

2 Thermodynamic phase transition of the 5-dimensional Reissner-Nordström-AdS black hole

Starting from the action

$$S = \frac{1}{16\pi G}\int d^{n+1}x\sqrt{-g}[R - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{n(n-1)}{l^2}], \tag{1}$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, and l is the AdS radius. We shall focus on the case of $n = 4$, in which the charged Reissner-Nordström-AdS black hole can be written as [9]

$$ds^2 = -f(r)dt^2 + f^{-1}(r)dr^2 + r^2[d\phi^2 + \sin^2\phi(d\theta^2 + \sin^2\theta d\psi^2)], \tag{2}$$

where $\phi \in (0,\pi), \theta \in (0,\pi), \psi \in (0,2\pi)$, are hyperspherical coordinates for the 3-sphere, and

$$f(r) = 1 + \frac{r^2}{l^2} - \frac{8M}{3\pi r^2} + \frac{4Q^2}{3\pi^2 r^4}, \tag{3}$$

with M and Q the mass and charge of the black hole. Whence we can get the Hawking temperature of this space time as

$$T = \frac{f'(r)}{4\pi}\mid_{r_+} = \frac{3\pi^2 r_+^6 - 8l^2\left(Q^2 - M\pi r_+^2\right)}{6l^2\pi^3 r_+^5}. \tag{4}$$

In addition, it follows from the Bekenstein-Hawking formula that the entropy of the black hole is given by

$$S = \frac{\pi^2 r_+^3}{2}, \tag{5}$$

where r_+ is the outer event horizon of the black hole, namely the largest root of the equation $f(r_+) = 0$. With this, the mass of the back hole can thus be expressed as the function of the event horizon

$$M = \frac{4l^2Q^2 + 3l^2\pi^2 r_+^4 + 3\pi^2 r_+^6}{8l^2\pi r_+^2}. \tag{6}$$

Substituting (5) and (6) into (4), we can get the relation between the temperature T and entropy S of the 5-dimensional Reissner-Nordström-AdS black hole

$$T = \frac{12S^2 + l^2\left(-2\pi^2Q^2 + 32^{1/3}\pi^{4/3}S^{4/3}\right)}{62^{2/3}l^2\pi^{5/3}S^{5/3}}. \tag{7}$$

In addition, with the relation $F = M - TS$, the Helmholtz free energy can be expressed as

$$F = \frac{5Q^2}{6\pi r_+^2} - \frac{1}{8}\pi r_+^2\left(-1 + r_+^2\right). \tag{8}$$

Note that this formula for our free energy has implicitly chosen the pure AdS as the reference spacetime because the free energy vanishes for pure AdS by this formula. Now

let us review the relevant phase transitions in the fixed charge ensemble by (7) and (8) in the $T-S$ plane.

To achieve this, we should first find the critical charge by the following equations

$$(\frac{\partial T}{\partial S})_Q = (\frac{\partial^2 T}{\partial S^2})_Q = 0. \tag{9}$$

Inserting (7) into (9), we can get the values for the critical charge and critical entropy

$$Q_c = \frac{l^2\pi}{6\sqrt{5}}, \tag{10}$$

$$S_c = \frac{l^3\pi^2}{6\sqrt{3}}. \tag{11}$$

Substituting these critical values into (7), we can get the critical temperature

$$T_c = \frac{\sqrt{3}(3+5)}{10l\pi}. \tag{12}$$

We plot the isocharge curves for different charges in Figure 1. For the case $Q = 0$, There is a minimum temperature $T_0 = \frac{\sqrt{2}}{\pi}$ [30], which is indicated by the red solid line in (a). When the temperature is lower than T_0, we have only a thermal AdS. When the temperature is higher than T_0, there are two additional black hole branches. The small branch is unstable while the large branch is stable. This can be justified by check the corresponding heat capacities, which is related to their slopes. The Hawking-Page phase transition occurs at the temperature given by $T_1 = \frac{3}{2\pi}$ [30], which is higher than T_0 and indicated by the red dashed line. This can be observed by the $F-T$ relation in (a) of Figure 2, where T_0 is the horizontal coordinate of the cusp and T_1 is the horizontal coordinate for the intersection of the stable branch and the horizontal axis. Obviously, when the temperature is lower than T_1, the thermal AdS is the most stable state. While when the temperature is higher than T_1, the most stable state is taken over by the large black hole branch.

For the case $Q \neq 0$, the phase structure is similar to that of the Van der Waals phase transition. That is, for a small charge, there is an unstable black hole interpolating between the stable small hole and stable large hole. The small stable hole will jump to the large stable hole at the critical temperature temperature $T_\star$, which is labeled by the red dashed line in (b) of Figure 1. As the charge increases to the critical charge, the small hole and the large hole merge into one and squeeze out the unstable phase. So there is an inflection point in (c) of Figure 1. The heat capacity is divergent in this case, the phase transition is therefore second order. As the charge exceeds the critical charge, we simply have one stable black hole at each temperature, which can be justified by the slope of the curve in (d) of Figure 1. The Van der Waals-like phase transition can also be observed from the $F-T$ relation. From (b) of Figure 2, we see a swallowtail structure, which corresponds to the unstable phase in (b) of Figure 1. The critical temperature $T_\star = 0.4526$ for the phase



(a) $Q = 0$ (b) $Q = \frac{\pi-1}{6\sqrt{5}}$

(c) $Q = \frac{\pi}{6\sqrt{5}}$ (d) $Q = \frac{\pi+1}{6\sqrt{5}}$

Figure 1: Relation between the entropy and temperature for different charges in the fixed charge ensemble. The red solid line corresponds to the minimum temperature of space time and the red dashed lines in (a), (b), (c) correspond to the locations of Hawking-Page phase transition, first order phase transition, and second order phase transition individually.



(a) $Q = 0$ (b) $Q = \frac{\pi-1}{6\sqrt{5}}$ (c) $Q = \frac{\pi}{6\sqrt{5}}$

Figure 2: Relation between the free energy and temperature for different charges. The horizontal coordinates of the red dashed lines correspond to the temperatures of the Hawking-Page phase transition, first order phase transition, and second order phase transition.

transition is apparently read off by the horizontal coordinate of the junction between the small black hole and the large black hole. As the temperature is lower than the critical temperature $T_\star$, the free energy of the small black hole is lowest, so the small hole is stable. As the temperature is higher than $T_\star$, the free energy of the large black hole is lowest, so the large hole dominates thereafter. The non-smoothness of the junction indicates that the phase transition is first order. When the charge is arriving at the critical charge Q_c, the swallowtail structure in (b) of Figure 2 shrinks into a point as is shown in (c) of Figure 2. The horizontal coordinate of the inflection point corresponds to the critical temperature T_c of the second order phase transition, which is consistent with the analytical result in (12).

For the first order phase transition in (b) of Figure 1, we would like to check whether Maxwell's equal area law holds with the following formula

$$A_1 \equiv \int_{S_1}^{S_3} T(S,Q)dS = T_\star(S_3 - S_1) \equiv A_3, \tag{13}$$

in which $T(S,Q)$ is defined in (7), S_1 and S_3 are the smallest and largest roots of the equation $T(S,Q) = T_\star$. After a simple calculation, we find $S_1 = 0.186987$ and $S_3 = 2.60575$. With these values, we find A_1 and A_3 in (13) equal 1.09481 and 1.09482 respectively. So the equal area law in the $T - S$ plane holds within our numerical accuracy.

For the second order phase transition in (c) of Figure 1, we are interested in the critical exponent associated with the heat capacity

$$C_Q = T\frac{\partial S}{\partial T}\,|_Q\,. \tag{14}$$

Near the critical point, writing the entropy as $S = S_c + \delta$ and expanding the temperature in terms of small δ, we find

$$T - T_c = \frac{\left(220l^2\pi^2Q^2 - 212^{1/3}l^2\pi^{4/3}S^{4/3} + 30S^2\right)}{2432^{2/3}l^2\pi^{5/3}S^{14/3}}(S - S_c)^3, \tag{15}$$

in which we have used (9). In this case, (14) further implies $C_Q \sim (T - T_c)^{-2/3}$, namely the critical exponent is $-2/3$, which is the same as the one from the mean field theory. In addition, taking logarithm to (15), we have a linear relation

$$\log \mid T - T_c \mid = 3 \log \mid S - S_c \mid + constant, \tag{16}$$

with 3 the slope. In what follows, we will use this logarithm to check the critical exponent for the analogous heat capacities in the framework of holography.

It it noteworthy that by holography the whole phase structure described above is not only for the bulk black hole but also for the dual boundary system, where the thermal entropy is simply given by the black hole entropy, and so on so forth.

3 Phase transition in the framework of holography

In this section we shall investigate the phase structures of some non-local observables such as two point correlation function, Wilson loop, and entanglement entropy in the dual fled theory by holography to see whether they have the same phase structure as the thermal entropy.

3.1 Phase transition of two point correlation function

According to the AdS/CFT correspondence, if the conformal dimension Δ of scalar operator $\mathcal{O}$ of dual field theory is large enough, the equal time two point correlation function can be holographically approximated as [31]

$$\langle \mathcal{O}(t_0, x_i)\mathcal{O}(t_0, x_j)\rangle \approx e^{-\Delta L}, \tag{17}$$

where L is the length of the bulk geodesic between the points (t_0, x_i) and (t_0, x_j) on the AdS boundary. Taking into account the spherical symmetry of the 5-dimensional Reissner-Nordström-AdS black hole, we can simply choose $(\phi = \frac{\pi}{2}, \theta = \theta_0, \psi = 0)$ and $(\phi = \frac{\pi}{2}, \theta = \theta_0, \psi = \pi)$ as the two boundary points. Then with θ to parameterize the trajectory, the proper length is given by

$$L = 2\int_0^{\theta_0} \mathcal{L}(r(\theta), \theta)d\theta, \quad \mathcal{L} = \sqrt{\frac{\dot{r}^2}{f(r)} + r^2}, \tag{18}$$

in which $\dot{r} = dr/d\theta$. Imagining θ as time, and treating $\mathcal{L}$ as the Lagrangian, one can get the equation of motion for $r(\theta)$ by making use of the Euler-Lagrange equation, that is

$$0 = \dot{r}^2 f'(r) - 2f(r)\ddot{r} + 2rf(r)^2, \tag{19}$$

which can be solved by imposing the following boundary conditions

$$\dot{r}(0) = 0, r(0) = r_0. \tag{20}$$

To explore whether the size of the boundary region affects the later phase structure, we here choose $\theta_0 = 0.14, 0.2$ as two examples. Note that for a fixed θ_0, the geodesic length is divergent, so it should be regularized by subtracting off the geodesic length in pure AdS with the same boundary region, denoted by L_0. To achieve this, we are required to set a UV cutoff for each case, which is chosen to be $r(0.139)$ and $r(0.199)$, respectively for our two examples. In this paper, we obtain L_0 also by numerics though there is an analytical result for $r(\theta_0)$ for pure AdS in Einstein gravity. We label the regularized geodesic length as $\delta L \equiv L - L_0$.

We plot the relation between T and δL for different θ_0 in Figure 3 and Figure 4. As shown in Figure 3 and Figure 4, δL demonstrates a similar phase structure as the thermal entropy. Moreover, we find that the minimum temperature T_0 as well as Hawking-Page



(a) $Q = 0$

(b) $Q = \frac{\pi-1}{6\sqrt{5}}$



(c) $Q = \frac{\pi}{6\sqrt{5}}$



(d) $Q = \frac{\pi+1}{6\sqrt{5}}$

Figure 3: Relation between the geodesic length and temperature in the fixed charge ensemble for differen charges at $\theta_0 = 0.14$. The red solid line corresponds to the location of the minimum temperature T_0, the dashed lines in (a), (b), (c) correspond individually to the locations of Hawking-Page phase transition T_1, first order phase transition $T_\star$, and second order phase transition T_c.



(a) $Q = 0$ (b) $Q = \frac{\pi-1}{6\sqrt{5}}$



(c) $Q = \frac{\pi}{6\sqrt{5}}$ (d) $Q = \frac{\pi+1}{6\sqrt{5}}$

Figure 4: Relation between the geodesic length and temperature in the fixed charge ensemble for differen charges at $\theta_0 = 0.2$. The red solid line corresponds to the location of the minimum temperature T_0, the dashed lines in (a), (b), (c) correspond individually to the locations of Hawking-Page phase transition T_1, first order phase transition $T_\star$, and second order phase transition T_c.



(a) $\theta_0 = 0.14$ (b) $\theta_0 = 0.2$

Figure 5: Relation between $\log \mid T - T_c \mid$ and $\log \mid \delta L - \delta L_c \mid$ near the critical point of second order phase transition for different θ_0.

phase transition temperature T_1 in (a), the first order phase transition temperature $T_\star$ in (b), and second order phase transition temperature T_c in (c) are exactly the same as those in $T - S$ plane, which justifies our notation. To be more specific, it is easy to check T_0 by locating the position of local minimum. But in order to confirm $T_\star$ and T_c, we are required to examine the equal area law for the first order phase transition and obtain $-2/3$ as the critical exponent for the second order phase transition, which are documented as follows.

In the $\delta L - T$ plane, we define the equal area law as

$$A_1 \equiv \int_{\delta L_1}^{\delta L_3} T(\delta L) d\delta L = T_\star(\delta L_3 - \delta L_1) \equiv A_3, \tag{21}$$

in which $T(\delta L)$ is an Interpolating Function obtained from the numeric result, and δL_1, δL_3 are the smallest and largest roots of the equation $T(\delta L) = T_\star$. For the case $\theta_0 = 0.14$, we find $\delta L_1 = 0.0000556147$, $\delta L_3 = 0.000194497$. Substituting these values into (21), we find $A_1 = 0.0000628401$, $A_3 = 0.0000628583$. For the case $\theta_0 = 0.2$, after simple calculation, we find $A_1 = 0.0000108924$, $A_3 = 0.0000108875$. It is obvious that for different θ_0, A_1 and A_3 are equal within our numeric accuracy. Thus the equal area law also holds in the $\delta L - T$ plane.

In addition, in order to investigate the critical exponent for the analogous heat capacity of the geodesic length. we are interested in the logarithm of the quantities $T - T_c$, $\delta L - \delta L_c$, in which T_c is the critical temperature defined in (12), and L_c is obtained numerically by the equation $T(\delta L) = T_c$. We plot the relation between $\log \mid T - T_c \mid$ and $\log \mid \delta L - \delta L_c \mid$ for different θ_0 in Figure 5, where these straight lines can be fitted as

$$\log \mid T - T_c \mid = \begin{cases} 23.2318 + 3.06832 \log \mid \delta L - \delta L_c \mid, & \text{for } \theta_0 = 0.14, \\ 31.9841 + 3.00077 \log \mid \delta L - \delta L_c \mid, & \text{for } \theta_0 = 0.2. \end{cases} \tag{22}$$

It is obvious that the slope is about 3, which indicates that the critical exponent is $-2/3$ for the analogous heat capacity and the phase transition is also second order at T_c for the geodesic length.

3.2 Phase transition of Wilson loop

In this subsection, we are going to study the phase structure of the Wilson loop, which in the bulk corresponds to the minimal area surface by holography. Wilson loop operator is defined as a path ordered integral of gauge field over a closed contour, and its expectation value is approximated geometrically by the AdS/CFT correspondence as [32]

$$\langle W(C)\rangle \approx e^{-\frac{A_\Sigma}{2\pi\alpha'}}, \tag{23}$$

where C is the closed contour, Σ is the minimal bulk surface ending on C with A its minimal area, and α' is the Regge slope parameter. Next we choose the line with $\phi = \frac{\pi}{2}$ and $\theta = \theta_0$ as our loop. Then we can employ (θ, ψ) to parameterize the minimal area surface, which is invariant under the ψ-direction by our rotational symmetry. Thus the corresponding minimal area surface can be expressed as

$$A = 2\pi \int_0^{\theta_0} r \sin\theta \sqrt{\frac{\dot{r}^2}{f(r)} + r^2} d\theta, \tag{24}$$

in which $\dot{r} = dr/d\theta$. Making use of the Euler-Lagrange equation, one can get the equation of motion for $r(\theta)$. Then with the boundary conditions $r'(0) = 0$, $r(0) = r_0$, we can further get the numeric result of $r(\theta)$. Similar to the case of geodesic length, we choose $\theta_0 = 0.14, 0.2$ as two examples and the corresponding UV cutoffs are set to be $r(0.139)$, $r(0.199)$. We label the regularized minimal area surface as $\delta A \equiv A - A_0$, where A_0 is the minimal area in pure AdS with the same boundary region. We plot the relation between δA and T for different θ_0 in Figure 6 and Figure 7. Comparing Figure 6 with Figure 7, we find they are the same nearly besides the scale of the horizonal coordinate. In other words, θ_0 affects only the value but not the phase structure of minimal area surface in the $T - \delta A$ plane. The result tells us that the similar phase structure also shows up for the minimal surface area. Here we concentrate only on scrutinizing the equal area law for the first order phase transition and the critical exponent of the analogous heat capacity for the second order phase transition.

First, in the $\delta A - T$ plane, the equal area law can be similarly defined as

$$A_1 \equiv \int_{\delta A_1}^{\delta A_3} T(\delta A) d\delta A = T_\star(\delta A_3 - \delta A_1) \equiv A_3, \tag{25}$$

in which $T(\delta A)$ is an Interpolating Function obtained from our numeric result, and δA_1, δA_3 are the smallest and largest roots of the equation $T(\delta A) = T_\star$, respectively. As the same as that of the geodesic length, for a fixed θ_0, we first obtain δA_1, δA_3, and then substitute these values into (25) to produce A_1, A_3. The concrete values are listed in Table 1. Obviously, for both θ_0, A_1 and A_3 are equal within the reasonable numeric accuracy. The equal area law thus holds in the $\delta A - T$ plane, which reinforces the fact that the minimal surface area has the same first order phase transition behavior as that of the thermal entropy.



(a) $Q = 0$ (b) $Q = \frac{\pi-1}{6\sqrt{5}}$



(c) $Q = \frac{\pi}{6\sqrt{5}}$ (d) $Q = \frac{\pi+1}{6\sqrt{5}}$

Figure 6: Relation between the minimal area surface and temperature in the fixed charge ensemble for differen charges at $\theta_0 = 0.14$. The red solid line corresponds to the location of the minimum temperature T_0, the dashed lines in (a), (b), (c) correspond individually to the locations of Hawking-Page phase transition T_1, first order phase transition $T_\star$, and second order phase transition T_c.

$\theta_0 = 0.14$	$\theta_0 = 0.2$
$T_\star$ =0.4526	$T_\star$ =0.4526
δA_1=0.0000434217| δA_3=0.000114835	δA_1=0.0000374703| δA_3=0.000248923
A_1=0.0000320531| A_3=0.0000323218	A_1=0.0000957474| A_3=0.0000957035

Table 1: Check of the equal area law in the $T - \delta A$ plane for different θ_0.



(a) $Q = 0$ (b) $Q = \frac{\pi-1}{6\sqrt{5}}$

(c) $Q = \frac{\pi}{6\sqrt{5}}$ (d) $Q = \frac{\pi+1}{6\sqrt{5}}$

Figure 7: Relation between the minimal area surface and temperature in the fixed charge ensemble for differen charges at $\theta_0 = 0.2$. The red solid line corresponds to the location of the minimum temperature T_0, the dashed lines in (a), (b), (c) correspond individually to the locations of Hawking-Page phase transition T_1, first order phase transition $T_\star$, and second order phase transition T_c.



(a) $\theta_0 = 0.14$ (b) $\theta_0 = 0.2$

Figure 8: Relation between $\log \mid T - T_c \mid$ and $\log \mid \delta A - \delta A_c \mid$ near the critical point of second order phase transition for different θ_0.

Second, in order to check whether the minimal surface area also demonstrates the same second order phase transition as the thermal entropy, we would like to evaluate the critical exponent of the analogous heat capacity at the critical point in the $\delta A - T$ plane. To this end, we plot the relations between $\log \mid T - T_c \mid$ and $\log \mid \delta A - \delta A_c \mid$ in Figure 8. The numerical results for these curves can be fitted as

$$\log \mid T - T_c \mid = \begin{cases} 27.5226 + 3.04698 \log \mid \delta A - \delta A_c \mid, & \text{for } \theta_0 = 0.14, \\ 24.692 + 3.00462 \log \mid \delta A - \delta A_c \mid, & \text{for } \theta_0 = 0.2. \end{cases} \tag{26}$$

With 3 the slope, we can conclude that the minimal surface area also has the same second order phase transition as the thermal entropy.

3.3 Phase transition of entanglement entropy

Holographic entanglement entropy is another non-local observable, and it has been used extensively to probe the superconductivity phase transition besides the thermalization process recently [33, 34, 35, 36, 37, 38, 39, 40]. In this subsection, we intend to employ it to probe the phase structure of a 5-dimensional Reissner-Nordström-AdS black hole. According to the formula in [41, 42], holographic entanglement entropy can be given by the area A_Σ of a minimal surface Σ anchored on the boundary entangling surface $\partial\Sigma$, namely

$$S = \frac{A_\Sigma(t)}{4G}. \tag{27}$$

For simplicity, we choose $\phi = \phi_0$ as our entangling surface and employ (ϕ, θ, ψ) to parameterize the minimal surface. But with the symmetry of (2), (27) can be rewritten as

$$S = 4\pi \int_0^{\phi_0} r^2 \sin^2\phi \sqrt{\frac{\dot{r}^2}{f(r)} + r^2} d\phi \tag{28}$$

with $\dot{r} = dr/d\phi$. Similarly, we can solve the equation of motion for $r(\phi)$ numerically, and eventually obtain the regularized entanglement entropy δS. We plot the relation between



(a) $Q = 0$ (b) $Q = \frac{\pi-1}{6\sqrt{5}}$

(c) $Q = \frac{\pi}{6\sqrt{5}}$ (d) $Q = \frac{\pi+1}{6\sqrt{5}}$

Figure 9: Relation between the entanglement entropy and temperature in the fixed charge ensemble for differen charges at $\phi_0 = 0.14$. The red solid line corresponds to the location of the minimum temperature T_0, the dashed lines in (a), (b), (c) correspond individually to the locations of Hawking-Page phase transition T_1, first order phase transition $T_\star$, and second order phase transition T_c.



(a) $Q = 0$ (b) $Q = \frac{\pi-1}{6\sqrt{5}}$

(c) $Q = \frac{\pi}{6\sqrt{5}}$ (d) $Q = \frac{\pi+1}{6\sqrt{5}}$

Figure 10: Relation between the entanglement entropy and temperature in the fixed charge ensemble for differen charges at $\phi_0 = 0.2$. The red solid line corresponds to the location of the minimum temperature T_0, the dashed lines in (a), (b), (c) correspond individually to the locations of Hawking-Page phase transition T_1, first order phase transition $T_\star$, and second order phase transition T_c.

$\phi_0 = 0.14$	$\phi_0 = 0.2$
$T_\star$ =0.4526	$T_\star$ =0.4526
δS_1=0.0000155355| δS_3=0.000123195	δS_1=0.000186918| δS_3=0.000517552
A_1=0.0000487431| A_3=0.0000487266	A_1=0.000148384| A_3=0.000149645

Table 2: Check of the equal area law in the $T - \delta S$ plane for different ϕ_0.



Figure 11: Relation between $\log \mid T - T_c \mid$ and $\log \mid \delta S - \delta S_c \mid$ near the critical point of second order phase transition for different ϕ_0

δS and T for $\phi_0 = 0.14, 0.2$ in Figure 9 and Figure 10, respectively. As one can see, it exhibits a similar behavior as the thermal entropy. To be more precise, we would like to check the equal area law with the following equation

$$A_1 \equiv \int_{\delta S_1}^{\delta S_3} T(\delta S) d\delta S = T_\star(\delta S_3 - \delta S_1) \equiv A_3, \tag{29}$$

in which $T(\delta S)$ is an Interpolating Function obtained from the numeric result and δS_1, δS_3 are the smallest and largest roots of the equation $T(\delta S) = T_\star$. For different $\phi_0 s$, the results of δS_1, δS_3 and A_1, A_3 are listed in Table 2. It is obvious that A_1 nearly equals A_3 regardless of the choice of ϕ_0. That is, the equal area law is also valid for the entanglement entropy.

To get the critical exponent of second order phase transition of entanglement entropy, we should find the slope of a linear function represented by $\log \mid T - T_c \mid$ and $\log \mid \delta S - \delta S_c \mid$, in which S_c is the critical entropy obtained numerically by the equation $T(\delta S) = T_c$. The numeric results for different ϕ_0 are plotted in Figure 11. The results for these curves can be further fitted as

$$\log \mid T - T_c \mid = \begin{cases} 26.653 + 3.00107 \log \mid \delta S - \delta S_c \mid, & \text{for } \phi_0 = 0.14 \\ 21.2674 + 2.92789 \log \mid \delta S - \delta S_c \mid, & \text{for } \phi_0 = 0.2 \end{cases} \tag{30}$$

One can see that the slope is always about 3 for different ϕ_0. So we can conclude that the entanglement entropy also has the same second order phase transition as the thermal entropy.

4 Concluding remarks

Investigation on the phase transition of the black holes is important and necessary. On one hand, it is helpful for us to understand the structure and nature of the space time. On the other hand, it may uncover some phase transitions of the realistic physics in the conformal field theory according to the AdS/CFT correspondence. It is well-known now that the Hawking-Page phase transition in the gravity system is dual to the confinement/deconfinement phase transition, and the phase transition of a scalar field is dual to the superconductivity phase transition in the dual conformal field theory.

In this paper, we investigated the van der Waals-like phase transition in the framework of holography so that we can explore whether there is a realistic similar phase transition in physics. Taking the 5-dimensional Reissner-Nordström-AdS black hole as the gravity background, we investigated the phase structure of the two point correlation function, Wilson loop, and holographic entanglement entropy. For all the non-local observables, we observed that the black hole undergos a van der Waals-like phase transition. This conclusion is reinforced by the investigation of the equal area law and critical exponent of the analogous heat capacity in which we found that the equal area law is valid always and the critical exponent of the heat capacity coincides with that of the mean field theory regardless of the size of the boundary region. In addition, we found the black hole undergos a Hawking-Page phase transition before the van der Waals-like phase transition for all the non-local observables. We also obtained the minimum temperature and Hawking-Page phase transition temperature. Our investigation thus provides a complete picture depicting the phase transition of charged AdS black hole in the framework of holography.

Acknowledgements

We would like to thank Rong-Gen Cai for his discussions. This work is supported by the National Natural Science Foundation of China (Grant Nos. 11405016, 11575270), China Postdoctoral Science Foundation (Grant No. 2016M590138), Natural Science Foundation of Education Committee of Chongqing (Grant No. KJ1500530), and Basic Research Project of Science and Technology Committee of Chongqing(Grant No. cstc2016jcyja0364).

References

[1] S. W. Hawking, *Particle creation by black holes,* Commun. Math. Phys. 43, 199 (1975)

[2] J. M. Bardeen, B. Carter, S. W. Hawking, *The four laws of black hole mechanics,* Commun. Math. Phys. 31, 161 (1973)

[3] J. M. Maldacena, *Large N limit of superconformal field theories and supergravity*, Int. J. Theor. Phys. 38, 1113 (1999)

[4] E. Witten, *Anti-de Sitter space and holography*, Adv. Theor. Math. Phys. 2, 253 (1998), [hep-th/9802150]

[5] S. S. Gubser, I. R. Klebanov, A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, Phys. Lett. B 428, 105 (1998), [hep-th/9802109]

[6] S. W. Hawking and D. N. Page, *Thermodynamics Of Black Holes In Anti-De Sitter Space*, Commun. Math. Phys. 87, 577 (1983)

[7] R. G. Cai, L. M. Cao, Y. W. Sun, *Hawking-Page Phase Transition of black Dp-branes and R-charged black holes with an IR Cutoff*, JHEP 0711, 039 (2007), [arXiv:0709.3568 [hep-th]]

[8] E. Witten, *Anti-de Sitter space, thermal phase transition, and confinement in gauge theories*, Adv. Theor. Math. Phys. 2, 505 (1998) [arXiv:hep-th/9803131]

[9] A. Chamblin, R. Emparan, C. V. Johnson and R. C. Myers, *Charged AdS black holes and catastrophic holography*, Phys. Rev. D 60, 064018 (1999) [hep-th/9902170]

[10] D. Kubiznak and R. B. Mann, *P-V criticality of charged AdS black holes*, JHEP 1207, 033 (2012), [arXiv:1205.0559 [hep-th]]

[11] J. Xu, L. M. Cao, Y. P. Hu, *P-V criticality in the extended phase space of black holes in massive gravity*, Phys. Rev. D 91, 124033 (2015), [arXiv:1506.03578 [gr-qc]]

[12] R. G. Cai, L. M. Cao, L. Li, R. Q. Yang, *P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space*, JHEP 09, 005 (2013), [arXiv:1306.6233 [gr-qc]]

[13] S. H. Hendi, A. Sheykhi, S. Panahiyan, B. Eslam Panah, *Phase transition and thermodynamic geometry of Einstein-Maxwell-dilaton black holes*, Phys. Rev. D 92, 064028 (2015), [arXiv:1509.08593 [hep-th]]

[14] R. A. Hennigar, W. G. Brenna, R. B. Mann, *P-V Criticality in Quasitopological Gravity*, JHEP 1507, 077 (2015), [arXiv:1505.05517 [hep-th]]

[15] S. W. Wei, Y. X. Liu, *Clapeyron equations and fitting formula of the coexistence curve in the extended phase space of charged AdS black holes*, Phys. Rev. D 91, 044018 (2015) [arXiv:1411.5749 [hep-th]]

[16] J. X. Mo, W. B. Liu, *P-V Criticality of Topological Black Holes in Lovelock-Born-Infeld Gravity*, Eur. Phys. J. C 74, 2836 (2014) [arXiv:1401.0785 [hep-th]]

[17] C. Niu, Y. Tian, and X. N. Wu, *Critical Phenomena and Thermodynamic Geometry of RN-AdS Black Holes*, Phys. Rev. D 85, 024017 (2012), [arXiv:1104.3066 [hep-th]]

[18] R. G. Cai, *Gauss-Bonnet Black Holes in AdS Spaces*, Phys. Rev. D 65, 084014 (2002), [arXiv:hep-th/0109133]

[19] V. Balasubramanian *et al.*, *Thermalization of Strongly Coupled Field Theories*, Phys. Rev. Lett. 106, 191601 (2011), [arXiv:1012.4753 [hep-th]]

[20] V. Balasubramanian *et al.*, *Holographic Thermalization*, Phys. Rev. D 84, 026010 (2011), [arXiv:1103.2683 [hep-th]]

[21] D. Galante and M. Schvellinger, *Thermalization with a chemical potential from AdS spaces*, JHEP 1207, 096 (2012), [arXiv:1205.1548 [hep-th]]

[22] E. Caceres and A. Kundu, *Holographic Thermalization with Chemical Potential*, JHEP 1209, 055 (2012), [arXiv:1205.2354 [hep-th]]

[23] X. X. Zeng, B. W. Liu, *Holographic thermalization in Gauss-Bonnet gravity*, Phys. Lett. B 726, 481 (2013), [arXiv:1305.4841 [hep-th]]

[24] X. X. Zeng, X. M. Liu, B. W. Liu, *Holographic thermalization with a chemical potential in Gauss-Bonnet gravity*, JHEP 03, 031 (2014) [arXiv:1311.0718 [hep-th]]

[25] X. X. Zeng, D. Y. Chen, L. F. Li, *Holographic thermalization and gravitational collapse in the spacetime dominated by quintessence dark energy*, Phys. Rev. D 91, 046005 (2015), [arXiv:1408.6632 [hep-th]]

[26] C. V. Johnson, *Large N Phase Transitions, Finite Volume, and Entanglement Entropy*, JHEP 1403, 047 (2014), [arXiv:1306.4955 [hep-th]]

[27] E. Caceres, P. H. Nguyen, J. F. Pedraza, *Holographic entanglement entropy and the extended phase structure of STU black holes*, [arXiv:1507.06069 [hep-th]]

[28] P. H. Nguyen, *An equal area law for the van der Waals transition of holographic entanglement entropy*, [arXiv:1508.01955 [hep-th]]

[29] X. X. Zeng, H. Zhang, L. F. Li, *Phase transition of holographic entanglement entropy in massive gravity*, [arXiv:1511.00383 [gr-qc]]

[30] R. Banerjee, D. Roychowdhury, *Thermodynamics of phase transition in higher dimensional AdS black holes*, JHEP 11, 004 (2011), [arXiv:1109.2433 [gr-qc]]

[31] V. Balasubramanian and S. F. Ross, *Holographic particle detection*, Phys. Rev. D 61, 044007 (2000), [arXiv:hep-th/9906226]

[32] J. M. Maldacena, *Wilson loops in large N field theories*, Phys. Rev. Lett. 80, 4859 (1998), [arXiv:hep-th/9803002]

[33] T. Albash, C. V. Johnson, *Holographic Studies of Entanglement Entropy in Superconductors*, JHEP 1205, 079 (2012), [arXiv:1202.2605 [hep-th]]

[34] R. G. Cai, S. He, L. Li, Y. L. Zhang, *Holographic Entanglement Entropy in Insulator/Superconductor Transition*, JHEP 1207, 088 (2012), [arXiv:1203.6620 [hep-th]]

[35] R. G. Cai, L. Li, L. F. Li, R. K. Su, *Entanglement Entropy in Holographic P-Wave Superconductor/Insulator Model*, JHEP 1306, 063 (2013), [arXiv:1303.4828 [hep-th]]

[36] L. F. Li, R. G. Cai, L. Li, C. Shen, *Entanglement entropy in a holographic p-wave superconductor model*, Nucl. Phys. B 894, 15-28 (2015), [arXiv:1310.6239 [hep-th]]

[37] R. G. Cai, S. He, L. Li, Y. L. Zhang, Holographic Entanglement Entropy in Insulator/Superconductor Transition, JHEP 1207, 088 (2012), [arXiv:1203.6620 [hep-th]]

[38] X. Bai, B. H. Lee, L. Li, J. R. Sun, H. Q. Zhang, *Time Evolution of Entanglement Entropy in Quenched Holographic Superconductors*, JHEP 040, 66 (2015), [arXiv:1412.5500 [hep-th]]

[39] R. G. Cai, L. Li, L. F. Li, R. Q. Yang, *Introduction to Holographic Superconductor Models*, Sci China-Phys Mech Astron, 58, 060401 (2015), [arXiv:1502.00437 [hep-th]]

[40] Y. Ling, P. Liu, C. Niu, J. P. Wu, Z. Y. Xian, *Holographic Entanglement Entropy Close to Quantum Phase Transitions*, [arXiv:1502.03661 [hep-th]]

[41] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, Phys. Rev. Lett. 96, 181602 (2006) [hep-th/0603001].

[42] S. Ryu, T. Takayanagi, *Aspects of Holographic Entanglement Entropy*, JHEP 0608, 045 (2006), [hep-th/0605073]